

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 39/40, 39/395 // (A61K 39/40, 39:02) (A61K 39/395, 39:002)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/14212</b> <b>(43) International Publication Date:</b> 9 April 1998 (09.04.98)
<b>(21) International Application Number:</b> PCT/US97/17259 <b>(22) International Filing Date:</b> 26 September 1997 (26.09.97) <b>(30) Priority Data:</b> 60/027,084 30 September 1996 (30.09.96) US <b>(71) Applicant (for all designated States except US):</b> EMBREX, INC. [US/US]; P.O. Box 13989, Research Triangle Park, NC 27709-3989 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> THOMA, John, A. [US/US]; 1206 Crestwood Drive, Fayetteville, AR 72701 (US). HADDAD, Eid, E. [US/US]; 1302 Suterland Road, Cary, NC 27511 (US). WHITFILL, Craig, E. [US/US]; 2004 Buckingham Way, Apex, NC 27502 (US). AVAKIAN, Alan, P. [US/US]; 6505 Wrenwood Avenue, Raleigh, NC 27607 (US). <b>(74) Agents:</b> SIBLEY, Kenneth, D. et al.; Myers Bigel Sibley & Sajovec, P.O. Box 37428, Raleigh, NC 27627 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> METHOD OF PRODUCING ACTIVE IMMUNITY WITH VACCINE CONJUGATE  <b>(57) Abstract</b>  A method of producing active immunity against a bacterial or protozoal disease in a subject comprises administering to the subject a vaccine conjugate comprising a live bacterial or protozoa and a neutralizing factor bound to the live bacterial or protozoa. The neutralizing factor is selected from the group consisting of antibodies and antibody fragments. The live bacterial or protozoa is one capable of producing disease in the subject, and the antibody or antibody fragment is one capable of neutralizing the live bacteria or protozoa.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## METHOD OF PRODUCING ACTIVE IMMUNITY WITH VACCINE CONJUGATE

This application claims the benefit of U.S. Provisional Application No. 60/027,084 filed September 30, 1996.

Field of the Invention

The present invention relates to methods of producing active immunity  
5 against a bacterial or protozoal disease by administering subjects a vaccine conjugate, which conjugate is comprised of a live bacteria or protozoa and a neutralizing antibody or fragment thereof.

Background of the Invention

10 Methods of producing active immunity against a viral disease by administering a vaccine conjugate, the vaccine conjugate comprised of a live virus and a viral neutralizing antibody, are described in U.S. Patents Nos. 5,397,568 and 5,397,569 to Whitfill et al. These references are concerned with viral diseases only.

Methods of treating coccidiosis, a protozoan disease of both birds and  
15 mammals caused by various *Eimeria* species, are described in U.S. Patent No. 4,935,007 to Baffundo et al. and U.S. Patent No. 5,055,292 to McDonald et al. *In ovo* inoculation against coccidiosis is described in published PCT applications WO 96/40233 and 96/40234.

Summary of the Invention

20 The present invention provides a method of producing active immunity against a bacterial or protozoal disease in a subject, the method comprising administering to the subject a vaccine conjugate comprised of a live bacteria or protozoa and a neutralizing factor bound to the live bacteria or protozoa. The neutralizing factor is  
25 selected from the group consisting of antibodies and antibody fragments. The antibody or antibody fragment is one capable of neutralizing the live bacteria or protozoa. The vaccine conjugate is administered in an amount effective to produce an immune response to the live bacteria or protozoa in the subject.

- 2 -

Another aspect of the present invention is a vaccine preparation useful for producing active immunity against a bacterial or protozoal disease in a subject. The vaccine preparation is a pharmaceutically acceptable formulation which comprises a vaccine conjugate. The vaccine conjugate comprises a live bacteria or protozoa and a neutralizing factor bound to the live bacteria or protozoa. The neutralizing factor is selected from the group consisting of antibodies and antibody fragments. The antibody or antibody fragment is capable of neutralizing the live bacteria or protozoa. The vaccine conjugate is included in the pharmaceutically acceptable formulation in an amount effective to produce an immune response to the live bacteria or protozoa in the subject.

Another aspect of the present invention is an article of manufacture comprising a closed, pathogen-impermeable, container and a sterile vaccine formulation as described above enclosed within the container.

#### Brief Description of Drawings

**Figure 1** graphs the oocyst output in avians vaccinated with a vaccine conjugate comprising 500 *E. acervulina* oocysts complexed with either 2.5, 25 or 150 µl of polyclonal antibody, compared to a non-vaccinated control (cntrl) and a control vaccinated with oocysts but without antibody. Oocyst output is used as a measure of infectivity.

**Figure 2** graphs the oocyst output in avians vaccinated with a vaccine conjugate comprising 500 *E. acervulina* oocysts complexed with either 25 or 150 µl of polyclonal antibody, compared to a non-vaccinated control (cntrl) and a control vaccinated with oocysts but without antibody. Oocyst output is used as a measure of infectivity.

**Figure 3** graphs oocyst output after vaccination and low-dose *E. acervulina* challenge. Vaccination used a vaccine conjugate comprising 500 *E. acervulina* oocysts complexed with either 2.5, 25 or 150 µl of polyclonal antibody; controls were a non-vaccinated control (cntrl) and a control vaccinated with oocysts but without antibody (0).

**Figure 4** graphs weight gain (in grams) in birds after vaccination and a high-dose *E. acervulina* challenge. Vaccination used a vaccine conjugate comprising 500 *E. acervulina* oocysts complexed with either 25 or 150 µl of polyclonal antibody; controls were a non-vaccinated control (cntrl) and a control vaccinated with oocysts but without

- 3 -

antibody (0).

Figure 5 graphs lesion scores in birds after vaccination and a high-dose *E. acervulina* challenge. Vaccination used a vaccine conjugate comprising 500 *E. acervulina* oocysts complexed with either 25 or 150 µl of polyclonal antibody; controls were a non-vaccinated control (cntrl) and a control vaccinated with oocysts but without antibody (0).

#### Detailed Description of the Invention

The present invention provides a vaccine preparation comprising a live organism (bacteria or protozoa) complexed with neutralizing antibodies specific to that organism. The amount of complexes neutralizing antibodies is such that the organism remains capable of inducing an active immune response, while at the same time providing some degree of protection against the deleterious effects of the pathogen. While applicants do not wish to be held to any single theory, it is currently believed that the present vaccine complex results in some form of delayed release of the pathogenic organism.

The present vaccine complex is thought to delay or initially protect the vaccinated subject from the pathogenic effects of the vaccine organism. However, this delay or initial protection is only temporary (in contrast to what would be expected using a dead or inactivated vaccine organism). The vaccine organism in the complex does ultimately infect the subject, inducing an active immunity. The degree of delay will be dependent on the amount of antibody used, the particular vaccine organism, and the subject to be vaccinated. Such a delay in infection is important when vaccinating young subjects, particularly when large numbers of subjects are to be vaccinated. For example, it is easier and more cost-efficient to vaccinate chicks *in ovo* compared to vaccinating newly hatched chicks.

In a preferred embodiment of the present invention, the neutralizing factor is provided in an amount which delays the appearance of pathological changes associated with infection of the subject by the live vaccine organism. The "delay" is comparative; the pathological changes are delayed in comparison to those which would occur if the live vaccine organism were administered without complexed neutralizing factor.

Use of the present vaccine conjugates are safer than the use of the

- 4 -

unconjugated organism yet are capable of inducing a protective active immune response. The term "safe" is used herein to indicate that the benefits of vaccination outweigh any harm in the majority of individuals vaccinated.

Antibodies used in practicing the present invention are bacterial or  
5 protozoal neutralizing antibodies. Bacterial or protozoal neutralizing antibodies are those which combat the infectivity of a bacteria or protozoa in vivo if the bacteria or protozoa and the antibodies are allowed to react together for a sufficient time. The source of the bacterial or protozoal neutralizing antibody is not critical. They may originate from any animal, including birds (e.g., chicken, turkey) and mammals (e.g., rat, rabbit, goat, horse).  
10 The bacterial or protozoal neutralizing antibodies may be polyclonal or monoclonal in origin. See, e.g., D. Yelton and M. Scharff, 68 American Scientist 510 (1980). The antibodies may be chimeric. See, e.g., M. Walker et al., 26 Molecular Immunology 403 (1989).

Bacterial or protozoal neutralizing antibodies used in practicing the  
15 present invention may be immunoglobulins of any isotype, including IgM, IgG, IgA, IgD, and IgE immunoglobulins. IgG and IgM are more preferred, and IgG immunoglobulins (e.g., IgG1, IgG2, IgG3, IgG4) are most preferred.

Antibody fragments used in practicing the present invention are fragments  
of bacterial or protozoal neutralizing antibodies which retain the variable region binding  
20 site thereof. Exemplary are F(ab')<sub>2</sub> fragments, F(ab') fragments, and Fab fragments. See generally Immunology: Basic Processes, 95-97 (J. Bellanti Ed. 2d ed. 1985).

Antibodies or antibody fragments used in practicing the present invention  
may have additional elements joined thereto. For example, a microsphere or  
microparticle may be joined to the antibody or antibody fragment, as described in U.S.  
25 Patent No. 4,493,825 to Platt, the disclosure of which is incorporated herein by reference.

The present invention is particularly advantageously employed with  
bacteria or protozoa which would be pathogenic (i.e., capable of causing disease) in the  
subject being treated if not for their conjugation to the neutralizing factor. The  
pathogenicity of the bacteria or protozoa may be inherent in the bacteria or protozoa itself  
30 or due to the susceptibility of the subject to be treated (e.g., birds in ovo). In general,  
many pathogenic bacteria or protozoa have the positive effect of evoking active immunity  
in subjects infected therewith, and many attenuated vaccine strains of bacteria or protozoa

- 5 -

have the capability of causing at least some disease in subjects. Hence, the term "pathogenic," as used to describe bacteria or protozoa herein, means that the harm caused to subjects by administration of the bacteria or protozoa outweighs any benefit which would result therefrom. An "active" or "live" organism refers to one which is not killed. A "vaccine organism" refers to one which is used for the induction of protective immune response, even though negative side effects may occur (in such cases the benefit of the active immunity outweighs any negative side effects). It is preferred that the bacteria or protozoa be a live organism one capable of producing an active immune response thereto in the subject being treated.

The vaccine conjugate is included in the vaccine formulations in an amount per unit dose sufficient to evoke an active immune response to the bacteria or protozoa in the subject to be treated. The term "immune response," as used herein, means any level of protection from subsequent exposure to the bacteria or protozoa which is of some benefit in a population of subjects, whether in the form of decreased mortality, decreased lesion scores, improved feed conversion ratios, or the reduction of any other detrimental effect of the disease, regardless of whether the protection is partial or complete.

With respect to the degree of protection provided by the neutralizing factor, the quantity of the neutralizing factor administered in combination with the bacteria or protozoa in the vaccine need not be sufficient to provide complete protection from the bacteria or protozoa, as long as the detrimental response produced by the bacteria or protozoa is reduced to a level at which the benefits of the immune response produced outweigh any harm resulting from the infection.

The term "subjects," as used herein, is intended to include, among other things, both mammals and birds. Exemplary mammals include mice, rats, pigs, rabbits, sheep, ferrets, dogs, cats, cows, horses and primates, including man. The term "bird" is intended to include males or females of any avian species, but is primarily intended to encompass poultry which are commercially raised for eggs or meat. Accordingly, the term "bird" is particularly intended to encompass hens, cocks and drakes of chickens, turkeys, ducks, geese, quail and pheasant.

Bacteria that may be used in carrying out the present invention include, but are not limited to, *Actinobacillosis lignieresi*, *Actinomyces bovis*, *Aerobacter*

- 6 -

- aerogenes*, *Anaplasma marginale*, *Bacillus anthracis*, *Borrelia anserina*, *Brucella canis*, *Clostridium chauvoei*, *C. hemolyticum*, *C. novyi*, *C. perfringens*, *C. septicum*, *C. tetani*, *Corynebacterium equi*, *C. pyogenes*, *C. renale*, *Cowdria ruminantium*, *Dermatophilus congolensis*, *Erysipelothrix insidiosa*, *Escherichia coli*, *Fusiformis necrophorus*,
- 5 *Haemobartonella canis*, *Hemophilus* spp. *H. suis*, *Leptospira* spp., *Moraxella bovis*, *Mycoplasma* spp. *M. hyopneumoniae*, *Nanophyetus salmincola*, *Pasteurella anatipestifer*, *P. hemolytica*, *P. multocida*, *Salmonella abortus-ovis*, *Shigella equirulis*, *Staphylococcus aureus*, *S. hyicus*, *S. hyos*, *Streptococcus agalactiae*, *S. dysgalactiae*, *S. equi*, *S. uberis*, and *Vibrio fetus* (for the corresponding diseases, see Veterinary Pharmacology and
- 10 Therapeutics 5th Edition, pg 746 Table 50.2 (N. Booth and L. McDonald Eds., 1982)(Iowa State University Press); and *Corynebacterium diphtheriae*, *Mycobacterium bovis*, *M. leprae*, *M. tuberculosis*, *Nocardia asteroides*, *Bacillus anthracis*, *Clostridium botulinum*, *C. difficile*, *C. perfringens*, *C. tetani*, *Staphylococcus aureus*, *Streptococcus pneumoniae*, *S. pyogenes*, *Bordetella pertussis*, *Pseudomonas aeruginosa*, *Campylobacter*
- 15 *jejuni*, *Brucella* spp., *Francisella tularensis*, *Legionella pneumophila*, *Chlamydia psittaci*, *C. trachomatis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella typhi*, *S. typhimurium*, *Yersinia enterocolitica*, *Y. pestis*, *Vibrio cholerae*, *Haemophilus influenza*, *Mycoplasma pneumoniae*, *Neisseria gonorrhoeae*, *N. meningitidis*, *Coxiella burnetii*, *Rickettsia mooseri*, *R. prowazekii*, *R. rickettsii*, *R. tsutsugamushi*, *Borrelia* spp.,
- 20 *Leptospira interrogans*, *Treponema pallidum*, and *Listeria monocytogenes* (for the corresponding diseases see R. Stanier et al., The Microbial World, pg. 637-38 Table 32.3 (5th Edition 1986).

- Protozoa that may be used in carrying out the present invention include, but are not limited to, the coccidiosis-causing *Eimeria* species (*E. tenella*, *E. necatrix*, *E.*
- 25 *brunetti*, *E. acervulina*, *E. mivati*, and *E. maxima*), *Anaplasma marginale*, *Giardia* species (e.g., *Giardia lamblia*), *Babesia* species (e.g., *B. canis*, *B. gibsoni*, *B. equi*, *B. caballi*, *B. bigemina*, *B. argentina*, *B. divergens*, and *B. bovis*) *Trichomonas foetus*, *Entamoeba histolytica*, and *Balantidium coli*; *Plasmodium* species (e.g., *P. falciparum*, *P. malariae*, *P. vivax*, and *P. ovale*), *Leishmania* species (e.g., *L. donovani*, *L. braziliensis*,
- 30 *L. tropica*, and *L. mexicana*), *Trypanosoma* species e.g., *T. brucei* and *T. cruzi*), *Entamoeba histolytica*, *Trichomonas vaginalis*, *Toxoplasma gondii*, and *Pneumocystis carinii*. As used herein, an "avian protozoan" is one known to infect avians.



- 7 -

The organisms may be administered in any suitable form, including spores or cysts thereof. For example, infective coccidial organisms may be administered in the form of sporulated oocysts, sporozoites, and sporocysts.

The exact number of the organisms to be administered in the form of a conjugate is not critical except that the number must be effective to engender an immunological response by the animal. In general, depending on the organism administered, the site and manner of administration, the age and condition of the subject, etc., the number of the organisms will range from 1, 10, or 100 organisms up to 1,000, 10,000, 100,000 or 1 million organisms. Where the organisms are administered as a conjugate to birds *in ovo* (within eggs), the dosage may be from 50, 100, or 500 up to 2,000, 10,000, 20,000, 30,000, 50,000 or 100,000 organisms or more.

Subjects may be administered vaccines of the present invention by any suitable means. Exemplary are by oral administration, by intramuscular injection, by subcutaneous injection, by intravenous injection, by intraperitoneal injection, by eye drop or by nasal spray. When the subject to be treated is a bird, the bird may be a hatched bird, including a newly hatched (i.e., about the first three days after hatch), adolescent, and adult birds. Birds may be administered the vaccine in ovo, as described in U.S. Patent No. 4,458,630 to Sharma (the disclosure of this and all other patent references cited herein is to be incorporated herein by reference).

The in ovo administration of the vaccine involves the administration of the vaccine to eggs. Eggs administered the vaccine of the present invention are fertile eggs which are preferably in the fourth quarter of incubation. Chicken eggs are treated on about the fifteenth to nineteenth day of incubation, and are most preferably treated on about the eighteenth day of incubation (the eighteenth day of embryonic development). Turkey eggs are preferably treated on about the twenty-first to twenty-sixth day of incubation, and are most preferably treated on about the twenty-fifth day of incubation.

Eggs may be administered the vaccine of the invention by any means which transports the compound through the shell. The preferred method of administration is, however, by injection. The site of injection is preferably within either the region defined by the amnion, including the amniotic fluid and the embryo itself, in the yolk sac, or in the air cell. Most preferably, injection is made into the region defined by the amnion. By the beginning of the fourth quarter of incubation, the amnion is sufficiently

- 8 -

enlarged that penetration thereof is assured nearly all of the time when the injection is made from the center of the large end of the egg along the longitudinal axis.

The mechanism of egg injection is not critical, but it is preferred that the method not unduly damage the tissues and organs of the embryo or the extraembryonic membranes surrounding it so that the treatment will not decrease hatch rate. A hypodermic syringe fitted with a needle of about 18 to 22 gauge is suitable for the purpose. To inject into the air cell, the needle need only be inserted into the egg by about two millimeters. A one inch needle, when fully inserted from the center of the large end of the egg, will penetrate the shell, the outer and inner shell membranes enclosing the air cell, and the amnion. Depending on the precise stage of development and position of the embryo, a needle of this length will terminate either in the fluid above the chick or in the chick itself. A pilot hole may be punched or drilled through the shell prior to insertion of the needle to prevent damaging or dulling of the needle. If desired, the egg can be sealed with a substantially bacteria-impermeable sealing material such as wax or the like to prevent subsequent entry of undesirable bacteria.

It is envisioned that a high speed automated egg injection system for avian embryos will be particularly suitable for practicing the present invention. Numerous such devices are available, exemplary being those disclosed in U.S. Patent No. 4,681,063 to Hebrank and U.S. Patents Nos. 4,040,388, 4,469,047, and 4,593,646 to Miller. All such devices, as adapted for practicing the present invention, comprise an injector containing the vaccine described herein, with the injector positioned to inject an egg carried by the apparatus with the vaccine. Other features of the apparatus are discussed above. In addition, if desired, a sealing apparatus operatively associated with the injection apparatus may be provided for sealing the hole in the egg after injection thereof.

Preferred egg injection apparatus for practicing the present invention is disclosed in U.S. Patents Nos. 4,681,063 and 4,903,635 to Hebrank, the disclosures of which are incorporated herein by reference. This device comprises an injection apparatus for delivering fluid substances into a plurality of eggs and suction apparatus which simultaneously engages and lifts a plurality of individual eggs from their upwardly facing portions and cooperates with the injection means for injecting the eggs while the eggs are engaged by the suction apparatus. The features of this apparatus may be combined with

- 9 -

the features of the apparatus described above for practicing the present invention.

Preferred subjects for carrying out the present invention are birds.

The method of the present invention is preferably carried out on birds in ovo.

5           A vaccine conjugate of the present invention is made by mixing the neutralizing factor with a live bacteria or protozoa in a pharmaceutically acceptable carrier for a time sufficient to form a live bacteria or protozoa-neutralizing factor conjugate (for example, by combining the neutralizing factor and bacteria or protozoa in a common liquid carrier prior to administration to a subject, until a conjugate is formed).

10       This can advantageously be carried out by simply adding hyperimmune sera containing neutralizing antibodies to an aqueous solution containing the live bacteria or protozoa. Vaccine formulations of the present invention preferably comprise the vaccine conjugate in lyophilized form or the vaccine conjugate in a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are preferably liquid, particularly aqueous, carriers.

15       For the purpose of preparing such vaccine formulations, the neutralizing factor and bacteria or protozoa may be mixed in sodium phosphate-buffered saline (pH 7.4), conventional media such as MEM, or bacteria growth medium. The vaccine formulation may be stored in a sterile glass container sealed with a rubber stopper through which liquids may be injected and formulation withdrawn by syringe.

20           The vaccine conjugate or complex of the present invention is a complex or conjugate of antibodies and live vaccine organisms; the bond between antibody and vaccine organism is a releasable bond and is not a covalent bond. The amount of neutralizing antibodies suitable for use with a given vaccine organism and a given subject can be readily determined using techniques available in the art. Use of too little antibody

25       will result in undesirably early or severe pathogenic effects caused by the vaccine organism; use of too much antibody may inactivate the vaccine organism completely or render it incapable of inducing a protective immune response.

          Vaccine formulations of the present invention may optionally contain one or more adjuvants. Any suitable adjuvant can be used, including chemical and

30       polypeptide immunostimulants which enhance the immune system's response to antigens. Preferably, adjuvants such as aluminum hydroxide, aluminum phosphate, plant and animal oils, and the like are administered with the vaccine conjugate in an amount

- 10 -

sufficient to enhance the immune response of the subject to the vaccine conjugate. The amount of adjuvant added to the vaccine conjugate will vary depending on the nature of the adjuvant, generally ranging from about 0.1 to about 100 times the weight of the bacteria or protozoa, preferably from about 1 to about 10 times the weight of the bacteria or protozoa.

The vaccine formulations of the present invention may optionally contain one or more stabilizer. Any suitable stabilizer can be used, including carbohydrates such as sorbitol, manitol, starch, sucrose, dextrin, or glucose; proteins such as albumin or casein; and buffers such as alkaline metal phosphate and the like. The use of a stabilizer is particularly advantageous when the vaccine formulation is a lyophilized formulation.

The present invention is explained in greater detail in the following non-limiting Examples.

- 11 -

## EXAMPLE 1

Bacterial Species

The bacterium *Pasteurella multocida* causes an acute highly contagious disease in many avian species. The disease, Fowl cholera, often occurs as a septicemic disease resulting in high morbidity and mortality. There are several live vaccines for Fowl cholera that can be administered to both chickens and turkeys.

A strain of *P. multocida* is complexed *in vitro* with antibodies specific to *P. multocida* to form bacterium-antibody complexes. Different ratios of bacterium to antibody are tested to determine a ratio that does not completely inactivate the bacterium, and still allows an active immune response to occur. These complexes are used as a vaccine in either chickens or turkeys. The responses of the vaccinee are followed and compared to the responses of birds vaccinated with the same dose of *P. multocida* vaccine not complexed with antibody. Lesions following vaccination, antibody response over time and general bird health are monitored throughout the trial.

Other bacteria are also tested in a similar manner. These bacteria include, but are not limited to, *Mycoplasma gallisepticum* in chickens or turkeys, *Bordetella avium* in turkeys, *Salmonella* species in chickens or turkeys, and *Salmonella* and *Listeria* species in rodents.

Vaccine conjugates are administered to birds either *in ovo* as described above or after hatch.

## EXAMPLE 2

Protozoal Species

The protozoan *Eimeria acervulina* (specifically, sporocysts and/or oocysts thereof) is complexed *in vitro* to antibodies specific to *Eimeria acervulina* to form protozoan-antibody complexes. These complexes are used as vaccines in chickens. The responses of the protozoan-antibody complex vaccinees are followed and compared to the responses of birds vaccinated with the same dose of *E. acervulina* not complexed with antibody. Different ratios of protozoan to antibody are tested to determine a ratio that does not completely inactivate the protozoan, and still allows an active immune response to occur. Oocyst output in vaccinee feces, intestinal absorptive ability (assessed by carotenoid uptake) and body weight are determined post vaccination. At 10-21 days post

- 12 -

vaccination birds in each vaccinated group are given a virulent challenge with *E. acervulina*. Oocyst output in vaccinates feces, body weight gain during the challenge period and intestinal absorptive ability (assessed by carotenoid uptake) are determined for vaccinates and non-vaccinated controls.

- 5                    Other *Eimeria* species are also tested in chickens, turkeys or rodents, as well as *Cryptosporidium* species in chickens or turkeys and *Histomonas meleagridis* in chickens or turkeys.

Vaccine conjugates are administered to birds either *in ovo* as described above or after hatch.

- 13 -

## EXAMPLE 3

Eimeria Oocyst Output Following Vaccination

Chickens were vaccinated and studied to determine the effects of complexing an *E. acervulina* oocyst vaccine with antibody.

5 Four vaccination strategies were studied. Treatment groups were vaccinated (by oral gavage) with 500 *E. acervulina* sporulated oocysts complexed with either 0, 2.5, 25 or 150 units of polyclonal antibody specific for *E. acervulina*. Each treatment was provided to three groups of five Leghorn chickens (15 birds total in each treatment group; 60 treatment birds overall). A control group of fifteen birds (5 birds in  
10 three repetitions) received no oocysts and no antibody, but were treated with oral gavage of 0.1 ml PBS administered on the day of hatch and were subsequently challenged.

Units of antibody is defined on a volumetric basis, where 1 $\mu$ l = 1 unit. An ELISA assay has been used to determine the "titer units" of the antibody preparation used in the present example; however, this assay has not been validated. By the ELISA assay,  
15 the *E. acervulina* antibody preparation had a titer of 90,782 units/ml. The doses used herein provide relative comparisons; the appropriate amount of antibody to be complexed with a given organism will depend on the organism, the antibody preparation, and the intended subject. One skilled in the art, using techniques known in the art, would be able to determine appropriate organism:antibody ratios for a given usage.

20 The oocysts and antibodies were mixed together in PBS for a minimum of one hour prior to vaccination at room temperature. Vaccine complex was then stored at 4 C until administration. Chickens were vaccinated on the day of hatch.

Oocyst output following vaccination was determined by collecting feces on days 4 to 8 post vaccination, and counting oocyst in the feces. The mean and standard  
25 deviation of oocyst output was determined for each group. As shown in Table 1, use of 150 $\mu$ l of antibodies complexed with 500 oocysts significantly reduced oocyst output.

- 14 -

**Table 1**  
**Infectivity – Oocyst Output per Bird**

5

Vaccination Treatment	Average Oocysts/Bird (x 10 <sup>6</sup> )
<u>None</u>	
Mean	0.00
Standard Deviation	0.00
N	3 <sup>2</sup>
<u>0 µl Abs + oocysts</u>	
Mean	(a) <sup>1</sup> 11.65
Standard Deviation	2.91
N	3 <sup>2</sup>
<u>2.5 µl Abs + oocysts</u>	
Mean	(a) <sup>1</sup> 10.18
Standard Deviation	2.54
N	3 <sup>2</sup>
<u>25 µl Abs + oocysts</u>	
Mean	(a) <sup>1</sup> 10.61
Standard Deviation	4.28
N	3 <sup>2</sup>
<u>150 µl Abs + oocysts</u>	
Mean	(b) <sup>1</sup> 5.40
Standard Deviation	1.94
N	3 <sup>2</sup>

1. a,b: differed at the 0.15 level by LSD test. Significance set at the 0.20 level due to the small sample size. Controls were not included in the statistical model.

2. N = Three groups of five birds in each group.

10



- 15 -

## EXAMPLE 4

**Elmeria Oocyst Output Following Challenge**

Birds in the treatment groups described in Example 1 were then challenged on Day 13 posthatch with 250 oocysts of *E. acervulina* in PBS administered by oral gavage. Feces was collected on days four to eight post-challenge, and average oocyst output was determined as a percentage of the oocyst output of the control group (the statistical model included the control group). Results are provided in Table 2. Greater output of oocysts following challenge indicates less protection against the pathogen challenge.

-16-

**Table 2 -- Protection after Challenge**

Vaccination Treatment	Average Oocyst Output per Bird; Post-challenge ( $\times 10^6$ )	Average Oocysts (% of control) <sup>2</sup>
<u>Control</u> <sup>2</sup>	A <sup>1</sup>	A <sup>1</sup>
Mean	23.08	100.00
Standard Deviation	1.67	7.22
N	3	3
<u>0 <math>\mu</math>l Abs + oocysts</u>	C <sup>1</sup>	C <sup>1</sup>
Mean	5.20	22.54
Standard Deviation	0.72	3.13
N	3	3
<u>2.5 <math>\mu</math>l Abs + oocysts</u>	B <sup>1</sup>	C <sup>1</sup>
Mean	8.72	37.76
Standard Deviation	2.45	10.60
N	3	3
<u>25 <math>\mu</math>l Abs + oocysts</u>	B <sup>1</sup>	B <sup>1</sup>
Mean	9.76	42.29
Standard Deviation	1.78	7.70
N	3	3
<u>150 <math>\mu</math>l Abs + oocysts</u>	B <sup>1</sup>	B <sup>1</sup>
Mean	10.49	45.47
Standard Deviation	2.33	10.10
N	3	3

1. A,B,C: differed at the .05 level by SNK test.

2. Control = non-vaccinated, challenged birds.

- 17 -

## EXAMPLE 5

**Protection Following Challenge**

Data provided in Table 2 was re-analyzed without including the control group data in the statistical model. Results are provided in Table 3.

5

**Table 3**

Vaccination Treatment	Average Oocysts Post-challenge (x10 <sup>6</sup> )	Average Oocysts (% of control)
<u>Control</u> <sup>2</sup>		
Mean	23.08	100.00
Standard Deviation	1.67	7.22
N	3	3
<u>0 µl Abs + oocysts</u>		
Mean	B <sup>1</sup> 5.20	B <sup>1</sup> 22.54
Standard Deviation	0.72	3.13
N	3	3
<u>2.5 µl Abs + oocysts</u>		
Mean	AB <sup>1</sup> 8.72	AB <sup>1</sup> 37.76
Standard Deviation	2.45	10.60
N	3	3
<u>25 µl Abs + oocysts</u>		
Mean	A <sup>1</sup> 9.76	A <sup>1</sup> 42.29
Standard Deviation	1.78	7.70
N	3	3
<u>150 µl Abs + oocysts</u>		
Mean	A <sup>1</sup> 10.49	A <sup>1</sup> 45.47
Standard Deviation	2.33	10.10
N	3	3

1. A,B,C: differed at the .05 level by SNK test.

2. Control = non-vaccinated, challenged birds.

10

The results provided in Examples 1-3 indicate that use of 150µl antibody in conjunction with the vaccination dose of 500 *E. acervulina* oocysts resulted in either a

- 18 -

lessening of the pathogenic effects of the vaccination (compared to use of same vaccination with lesser amounts of antibody, or no antibody; indicated by decreased oocysts output after vaccination), or possibly a delay in the pathogenic effects of the vaccination dose. As shown in Table 1, the use of 150µl of antibody complexed to the vaccine oocysts resulted in a lower infectivity level than vaccination without antibodies or the use of lesser amounts of antibodies ( $p \leq 0.15$ ).

As shown in Table 2, birds vaccinated with antibody-oocyst vaccine had reduced oocyst output after challenge with 250 *E. acervulina* oocysts, compared to unvaccinated birds. These results indicate that the antibody-oocyst vaccine preparation is effective in inducing a protective immune response against the challenging pathogen. Protection was highest in the treatment group vaccinated with oocysts but without any antibody, with a significance level of 0.05. As shown in Table 3, among birds treated with antibody-oocyst vaccine, the highest protection is again seen in the group treated with oocyst but no antibody.

The above results indicate that use of a vaccine complex of antibody and oocyst resulted in either a lessening of the pathogenic effects of the vaccine oocysts, or a delay in the pathogenic effects of the vaccine oocysts, while still engendering a protective immune response. Either effect would be expected to increase the safety of a vaccine, either by allowing administration of the vaccine to subjects who are more susceptible to the pathogenic effects of the vaccine organism, or to subjects at a younger age (such as to avians *in ovo*).

The foregoing are illustrative of the present invention, and are not to be taken as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

#### EXAMPLE 6

##### Use of Antibody-Oocyst (*Eimeria acervulina*) Vaccine Conjugate in Low-Dose Challenge Model

This study tested a 500 *Eimeria acervulina* oocyst vaccine complexed with varying amounts of antibody (2.5 to 150µl). Treatments were compared to a non-vaccinated control, and a control vaccinated without antibodies. All treatments were

- 19 -

administered on the day of hatch. Oocyst output after vaccination on Days 4-8 was measured for all treatments. Oocyst output was also measured after a Day 13 low dose challenge.

5       **Materials and Methods:** Hyvac SPF leghorns were used to rule out any effect of maternal antibodies. Polyclonal antibody was produced from chickens immunized with *E. acervulina* oocysts. Two antibody preparations were combined to create an *E. acervulina* antibody with a final titer of 90,782. Treatment groups and experimental design are shown in Table 4.

10

Table 4

Treatment	Oocyst Dose	Antibody (μl)	# of Birds/Rep	# Reps	Total # of Birds
1 (A,B,C)	0	0	5	3	15
2 (A,B,C)	500	0	5	3	15
3 (A,B,C)	500	2.5	5	3	15
4 (A,B,C)	500	25	5	3	15
5 (A,B,C)	500	150	5	3	15

15       The vaccine complex was produced by mixing oocysts (USDA #12 Lot 28-131-36) with antibody in the appropriate volume. The complex was incubated at ambient temperature for one hour before administration. Birds were gavaged on Day of Hatch with a 200 μl dose of the respective treatment. Fecal material was collected from Day 4 to Day 8. Fecal samples were processed and counted using McMaster's chambers to  
20       determine oocyst output per bird.

Birds were moved to a brooder unit and challenged on Day 13 post-hatch with a low dose challenge (250 *E. acervulina* oocysts). Feces was collected Days 4-8 post challenge and enumerated as described above.

25

The vaccinated control showed  $\approx 12 \times 10^6$  oocyst output per bird. The 2.5μl and 25 μl antibody treatments showed similar results, however, the 150μl antibody treatment may have had an inhibitory effect on oocyst output, having only 46% of the

- 20 -

output compared to the control. See Figure 1.

After a low dose challenge, similar results were seen in all the antibody vaccinated treatment groups. Figure 3. The three antibody treatment groups averaged  
5 approximately 40% output of control. The vaccinated control exhibited an output of only 23% of control.

**EXAMPLE 7****Use of Antibody-Oocyst (*Eimeria acervulina*) Vaccine Conjugate in High-Dose****5 Challenge Model**

Two antibody-oocyst vaccine conjugate formulations were tested in a high-dose challenge model. Infectivity was measured by oocyst output and response to challenge  
 10 was measured by weight gain and lesion scores. The vaccine conjugates consisted of 500 *E. acervulina* oocysts complexed with either 25 or 150µl of antibody (as described in **Example 6**); see **Table 5**. The same bird strain, antibody, and oocyst lot was used as in **Example 6** above.

15

**Table 5**

Treatment #	Oocyst Dose	Antibody (µl)	# of Birds/ Rep	# Reps	Total # Of Birds
1 (A,B,C)	0	0	10	3	30
2 (A,B,C)	500	0	10	3	30
3 (A,B,C)	500	25	10	3	30
4 (A,B,C)	500	150	10	3	30

Vaccines were prepared as described above and administered on Day 0 post-hatch in 200 µl volume. Fecal material was collected from Day 4-8 and enumerated.  
 20

Post-challenge parameters measured in the present experiment differed from **Example 6**. A high dose challenge was administered to all treatment groups on Day 13 and weights of each individual bird were recorded. After eight days (Day 21) the birds  
 25 were weighed and the lesions scored.

Oocyst output was lower in this experiment for the vaccinated control, as well as for the two antibody treatments, compared to **Example 6**; the vaccinated control (no antibody) showed a 20-fold decrease in oocyst output (see **Figure 2**). The cause of this  
 30 reduction is not clear.

- 22 -

A high dose challenge (500 oocyst challenge) was administered to all the treatment groups and weight gain and lesion scores were examined. Weight gain results (Figure 4) did not show a difference among the treatment groups, including vaccinated and non-vaccinated controls. Lesion score data (Figure 5) indicated protection of all vaccinated groups over the non-vaccinated control.

## 10 EXAMPLE 8

### *Pasteurella multocida*

**Production of antiserum to *P. multocida*:** Ten SPF chickens were housed in a clean room from hatch; at four weeks of age each bird received (subcutaneous injection in the neck) 0.5 mls of Solvay's "Pabac", a commercial inactivated *P. multocida* oil emulsion vaccine containing serotypes 1, 3 and 4; at eight weeks of age, an additional 0.5 mls was injected subcutaneously into the neck and another 0.5 mls was injected intramuscularly in the right breast. Approximately 20 mls of blood was removed from each bird by cardiac puncture at ten weeks of age. Antiserum was collected from the blood, pooled, and filtered through a .45 µm filter. After numerous sterility tests all produced negative results, the antisera was placed into different sized vials and placed in a -20 degree C freezer until use.

**Isolation and titer determination of the Cu and M9 strains of *P. multocida*:**  
The Cu and M9 strains of *P. multocida* were grown from live vaccines, Choleramune Cu and Multimune M respectively, each produced by Biomune. It was determined that these strains of *P. multocida* freeze best at -70 degrees C in a mixture of 90% culture and 10% glycerol.

## 30 EXAMPLE 9

### **Hatchability of Eggs Inoculated at Day 18 with *P. multocida***

This experiment was designed to determine if different numbers of colony forming units (CFUs) of *P. multocida* strains Cu and M9 affect the hatchability of SPF



-23-

eggs following *in ovo* inoculation at day 18 of incubation. Each strain of *P. multocida* was diluted in Phosphate Buffered Saline (PBS) to produce three dilutions: 1000 CFUs per 0.1 ml; 100 CFUs per 0.1 ml; and 10 CFUs per 0.1 ml. At day 18 of incubation, 0.1 mls of each dilution for each strain was inoculated into fourteen SPF eggs. Thirteen

5 eggs were inoculated with 0.1 ml of a mixture of Brain Heart Infusion Broth (BHI), PBS, and Glycerol (vehicle control) and thirteen eggs received no inoculation.

The results in Table 6 show that hatchability of SPF eggs was severely depressed in groups 2 through 7.

10 Table 6

**Hatchability of SPF Eggs Following Inoculation at Day 18  
with *P. multocida***

	<u>Group 1</u> BHI, PBS Glycerol	<u>Group 2</u> Cu, 1000 CFUs	<u>Group 3</u> Cu, 100 CFUs	<u>Group 4</u> Cu, 10 CFUs	<u>Group 5</u> M9, 1000 CFUs	<u>Group 6</u> M9, 100 CFUs	<u>Group 7</u> M9, 10 CFUs	<u>Group 8</u> No Inocu- lation
Normal Hatched	12			3			1	11
Un-Hatched	1	14	14	11	14	14	13	2
% Hatched	92	0	0	21	0	0	7	85

15 Each inoculated egg was inoculated with 0.1 ml of a culture/PBS mixture containing the appropriate number of CFUs. The dilutions containing  $1 \times 10(2)$  CFUs/ml were titrated for each strain and the titers differed somewhat from the target numbers. Group 4 actually contained 10.5 CFUs/egg, Group 3 contained 105 CFUs/egg, and Group 2 contained 1050 CFUs/egg. Group 7 contained 12.5 CFUs/egg,

20 Group 6 contained 125 CFUs/egg, and Group 5 contained 1250 CFUs/egg. Group 1 eggs were inoculated with 0.1 ml of a BHI/PBS/Glycerol mixture.

#### EXAMPLE 10

25 Colony Growth of *P. multocida* after Complexing with Chicken Antiserum

One ml of chicken *P. multocida* antiserum (see Example 8) was removed from -20 degrees C and thawed at room temperature. The serum was serially diluted, 2-fold,

- 24 -

in PBS. One half ml of each serum dilution was mixed with 0.5 ml of *P. multocida* Cu strain culture containing 100 CFUs/0.5 ml. This mixture reacted together for one hour at room temperature. When one hour had elapsed, 0.5 ml of each serum dilution mixture was plated on TSA plates in duplicate and the 200 CFUs/ml *P. multocida* Cu strain stock solution was plated in triplicate. The plates were incubated for 48 hours at 37 degrees C. Colonies were counted after 24 hours and 48 hours of incubation. Results are shown in Tables 7 and 8.

10 **Table 7**

Serum Dilution	24 Hours CFUs/0.5 ml of mixture	Avg	CFUs/1.0ml	#CFUs expected*	48 Hours change in CFU count/ 1.0 ml of mixture
1:16	114 + 96	= 105 x 2 =	210	152	no change
1:32	72 + 75	= 73.5 x 2 =	147	152	no change
1:64	92 + 86	= 89 x 2 =	178	152	no change
1:128	98 + 71	= 84.5 x 2 =	169	152	no change
1:256	81 + 92	= 86.5 x 2 =	173	152	no change
1:512	64 + 80	= 72 x 2 =	144	152	no change
1:1024	63 + 101	= 82 x 2 =	164	152	no change
1:2048	73 + 73	= 73 x 2 =	146	152	no change
1:4096	68 + 81	= 74.5 x 2 =	149	152	no change
1:8192	60 + 61	= 60.5 x 2 =	121	152	no change
1:16384	42 + 61	= 51.5 x 2 =	103	152	+ 1 colony
1:32768	56 + 59	= 57.5 x 2 =	115	152	no change

\* This is the number of CFUs expected in 1 ml of mixture, from the Table 8 titer.

15 **Table 8**  
**Titer of the 2 x 10<sup>2</sup> Stock Solution of *P. multocida* Cu Strain**

CFU/0.5 ml	Average Per 1.0 ml
176	152 x 2 = 304
131	
150	

20 Stock solution samples in Table 8 were plated after all mixtures were prepared. The solution did not sit for an additional hour. A mixture of 0.5 ml of this stock solution and 0.5ml of antiserum yields an expected 152 CFUs/ml of mixture.

- 25 -

The results show a decrease in the number of bacterial CFUs from the least dilute antiserum to the most dilute antiserum (Table 7). This may be due to reaction time. The most dilute antiserum reacted longer with the live culture than the least  
5 dilute antiserum (20-30 minutes longer). The increased time may have lead to greater numbers of colonies dying off. Repeated vortexing of the stock solution (3-4 CFUs/ml) may have also contributed to the observed decrease in colony count. The next Example investigates this trend further.

10

**EXAMPLE 11**

This experiment investigated the reaction between colonies of *P. multocida* Cu strain and the same dilutions of the *P. multocida* antiserum as shown in Table 6. Instead of vortexing the "2x10(2)" stock solution of *P. multocida* Cu strain continually,  
15 as in the previous experiment, the solution was gently mixed by pipetting. To form each antibody-bacterium complex, one ml of antiserum dilution was mixed with one ml of stock culture dilution. This mixture was allowed to react for four hours. The stock solution was plated on TSA plates in triplicate and was found to contain 186 CFUs/ml (Table 9). The remainder of this solution was also allowed to remain at room  
20 temperature for four hours.

Following the four hour reaction time, 0.5 ml of each antibody-bacterium mixture was mixed and plated in duplicate on TSA. The 186 CFU/ml stock solution was mixed and plated on TSA in triplicate (0.5 ml/plate). Colonies were counted at 24 hours after incubation at 37 degrees C and again after 96 hours. The results are  
25 provided in Tables 9 and 10.

- 26 -

Table 9

Titer of the planned  $2 \times 10^2$  CFUs/ml final stock solution of *P. multocida* strain Cu

5

Immediately after Adding 1 ml samples to serum			After sitting at room temperature for 4 hours		
CFU/0.5 ml	Average	Per 1 ml	CFU/0.5 ml	Average	Per 1 ml
93	93	186	39	44.7	89
			47		
			48		

Table 10

10

Serum Dilution	24 Hours CFUs/0.5 ml	Avg	CFUs/1.0ml	#CFUs expected*	96 Hours
1:16	114 + 92	103	206	93	no change
1:32	79 + 73	76	152	93	no change
1:64	50 + 48	49	98	93	no change
1:128	58 + 69	63.5	127	93	no change
1:256	52 + 67	59.5	119	93	no change
1:512	72 + 63	67.5	135	93	no change
1:1024	55 + 63	59	118	93	no change
1:2048	55 + 62	58.5	117	93	no change
1:4096	51 + 53	52	104	93	no change
1:8192	28 + 34	31	64	93	no change
1:16384	20 + 26	23	46	93	no change
1:32768	24 + 20	22	44	93	no change

The data in Table 10 shows the same decreasing colony count with increasing antiserum dilution as was seen in the previous experiment, slightly more pronounced.

- 15 There appears to be more CFUs/ml than expected in dilution mixtures 1:16 through 1:4096. Similar findings were also presented in Table 7. The colony counts found in Table 9 suggest the CFUs die off in PBS over time in the absence of serum. This may account for the lower than expected colony counts in dilutions 1:8192 through 1:32768 in both studies. The antiserum may be demonstrating some growth inhibiting
- 20 capabilities at the higher dilutions while exhibiting growth enhancement effects in the least dilute mixtures. The next experiment was conducted to investigate possible

-27-

causes of these observations.

#### EXAMPLE 12

5           The effects of *P. multocida* antiserum and negative serum (chicken serum containing no *P. multocida* antibodies) on the growth of the M9 strain of *P. multocida* was compared *in vitro*. Serum dilutions and *P. multocida* culture dilutions were prepared as in Example 11. One ml of culture containing 139 CFUs per ml (Table 6) was mixed with one ml of dilute serum. Three dilutions (1:16, 1:512, and  
10 1:32768) of negative serum were prepared and mixed with the appropriate amount of bacterial culture. The bacterial culture/serum dilution mixtures reacted together for one hour at room temperature.

          The serum antibody-bacterium mixtures were plated on TSA plates (0.5 ml) in duplicate and the negative serum bacterium mixtures were plated in triplicate following  
15 the one hour reaction time. The *P. multocida* strain M9 stock solution was plated in triplicate after adding 1.0 ml to all antiserum and negative serum dilutions and again after allowing it to remain at room temperature for one hour. Colonies were counted after incubating for 24 hours at 37 degrees C and any changes in counts were noted after 48 hours of incubation. The results of these counts are given in Tables 11 and 12.

- 28 -

Table 11

Titer of the planned 2 x 10(2) CFU/ml final stock solution of *P. multocida* M9

Immediately after adding 1 ml samples to serum			After sitting at room temperature for 1 hour		
CFU/0.5 ml	Average	Per 1 ml	CFU/0.5 ml	Average	Per 1 ml
68			61		
76	69.3	138.6	63	64.3	128.6
64			69		

5

Table 12

Serum Dilution	<i>P. multocida</i> antiserum after 24 hrs			# CFUs expected in 1 ml*	Negative antiserum after 24 hours		
	CFUs/ 0.5 ml	Avg.	Per 1 ml		CFUs/ 0.5 ml	Avg.	Per 1 ml
1:16	36+34	35	70	69	34+55+50	46.3	92.6
1:32	39+33	36	72	69			
1:64	40+37	38.5	77	69			
1:128	53+33	43	86	69			
1:256	33+35	34	68	69			
1:512	34+51	42.5	85	69	44+43+40	42.3	84.6
1:1024	36+35	35.5	71	69			
1:2048	32+37	34.5	69	69			
1:4096	34+29	31.5	63	69			
1:8192	34+44	39	78	69			
1:16384	24+25	24.5	49	69			
1:32768	18+38	28	56	69	40+39+33	37.3	74.6

\* Number of CFUs expected in 1 ml of mixture from Table 11 titer.

There were no changes in the colony counts after 48 hours of incubation at 37 degrees C.

10

The data in Examples 10-12 show initial increases leading to gradual decreases in the number of CFUs as antiserum to *P. multocida* became more dilute. The combined data suggest that both strains of *P. multocida* may actually use the serum as a growth medium. As the serum concentration decreases so does the growth of *P. multocida*. Some of the results suggest that *P. multocida* dies off when diluted in PBS and sits at room temperature for between one and four hours. This may account for the fewer than expected CFUs observed from the 1:8192 dilutions through 1:32768 dilutions. As the serum becomes less concentrated, the PBS concentration increases.

There was not a large difference, though, in the stock colony counts in the present

20

- 29 -

example, before and after a one hour waiting period, yet the low counts in the higher dilutions still existed.

The colony counts for the serum mixtures containing no antibodies to *P. multocida*, when compared to their *P. multocida* antibody-containing counterparts and stock counts, may be showing some growth inhibition capabilities of the antiserum with *P. multocida* antibodies. CFU counts in the samples without *P. multocida* antiserum started high and decreased but not to a level lower than expected when compared to the stock CFU counts after one hour. CFU counts were lower in two of three samples containing antibodies to *P. multocida* than in the negative serum samples without *P. multocida* antibodies. A growth inhibiting of *P. multocida* antiserum may be masked by other factors present.

### EXAMPLE 13

#### Hatchability of Eggs Injected at Day 18 of Incubation with *P. multocida* antiserum-*P. multocida* CFUs

This study was designed to test the effect of serum antibody-bacterium complexes when administered *in ovo* to SPF eggs. The same number of CFUs (five were targeted) of *P. multocida* strain M9 was mixed with varying amounts of *P. multocida* antiserum and then 0.1 ml of each mixture was inoculated into fifteen SPF eggs in each of seven groups. A 100 CFUs/ml stock solution was prepared using *P. multocida* strain M9 culture. To insure that each egg received the same number of CFUs, the appropriate amount of serum was mixed with the appropriate amount of the stock solution for each group at five minute intervals. These mixtures remained at room temperature for thirty minutes, after which time, 0.1 ml was inoculated into fifteen eggs at day eighteen of incubation. The 100 CFUs/ml stock solution was plated in triplicate following the preparation of the final group's mixture and again after the final group's mixture reacted for 30 minutes (Table 13). The plates were all incubated at 37 degrees C for 24 hours, after which, colonies were counted. Table 14 provides hatchability data for each group.

-30-

Table 13

	Titer of the planned $1 \times 10(2)$ <i>P. multocida</i> Strain M9 Stock Solution		
	CFUs/0.5ml	Average	Per 1 ml
Pre- 30 minutes	38 28 30	32	64
Post- 30 minutes	30 30 27	29	58



-31-

Table 14

Hatchability effects of inoculating SPF eggs at day 18 of incubation with 1 ml of mixtures containing antiserum to *P. multocida* and CFUs of the M9 strain of *P. multocida*.

	<u>Group 1</u> no serum +3.2CFUs	<u>Group 2</u> 0.1 µl serum +3.2 CFUs	<u>Group 3</u> 1.0µl serum +3.2CFUs	<u>Group 4</u> 10.0µl serum +3.2 CFUs	<u>Group 5</u> 25.0 µl serum +3.2 CFUs	<u>Group 6</u> 50.0µl serum +3.2 CFUs	<u>Group 7</u> 50.0µl serum 50.0µl PBS/ egg
Normal Hatched		2	1		2	4	15
Hatched (Dead)							
Hatched (clinically affected)	1		3	3	3	5	
Un-hatched	14	13	11	12	10	6	
% hatched	6.7	13.3	26.7	20.0	33	60.0	100

This study was originally designed to inoculate each egg that was to receive bacteria with 5.0 CFUs mixed with the appropriate volume of *P. multocida* antiserum. The titer information found in Table 13 shows that the eggs received a number of CFUs closer to 3.2 than to 5 and that little colony loss occurred as a result of the 30 minute reaction period. The results in Table 14 show that even small numbers of CFUs of *P. multocida* strain M9 are devastating to SPF eggs when administered at day eighteen of incubation. The controls in group 7 experienced a 100% hatch. The hatch was severely affected in the other groups. Of these groups, 5 and 6 contained the next highest percentages of total hatched birds. The eggs in group 6 were inoculated with the highest ratio of antiserum to *P. multocida* (50µl + 3.2 CFUs) and experienced a 60% overall hatch with a 27% normal hatch. This trend suggests that antiserum to *P. multocida*, when combined with the live bacteria, may provide some degree of protection to a chicken embryo by either decreasing or delaying the pathogenic effects of the bacterium.

The data in Tables 12 and 14, comparing serum with and without *P. multocida* antibodies and different amounts of serum antibodies respectively, show a possible inhibitory effect of serum antibodies to *P. multocida* on the growth, and perhaps the

- 32 -

pathogenic effects, of the organism. In Table 14 it was shown that the two highest amounts of antibody (Groups 5 and 6) resulted in better hatches compared to the bacterium alone (Group 1).

5

**EXAMPLE 14****Growth of *M. gallisepticum*; Production of Hyperimmune Sera**

A culture of the bacterium *Mycoplasma gallisepticum* strain F was obtained from North Carolina State University, Mycoplasma Lab, College of Veterinary  
10 Medicine. The F strain of *M. gallisepticum* is used in the commercial layer industry as a live vaccine. Forty milliliters of Frey's Media supplemented with 15% swine serum (FMS) was inoculated with 1.33 mls of the bacterial culture. This mixture was then incubated for approximately 18 hours at 37 degrees C and the grown culture was mixed 80/20 with sterile glycerol for freezing at -70degrees C. Sterility of this mixture was  
15 tested on Trypticase Soy Agar (TSA) and no extraneous organisms grew. Titer determination for the *M. gallisepticum* strain F stock culture after 24 hours at -70degrees C was  $5.8 \times 10(8)$  CFUs/ml.

Antiserum to *M. gallisepticum* strain R was purchased from the NCSU Mycoplasma Lab. The antiserum was produced by hyperimmunizing New Zealand  
20 White rabbits with inactivated *M. gallisepticum* strain R in adjuvant. Rabbits were immunized by intramuscular and intradermal injections three times prior to blood collection. This antiserum is designated as MGA.

**EXAMPLE 15****25 Growth Inhibition Effect of MGA on *M. gallisepticum* Strain F**

This experiment investigated the growth of a given amount of *M. gallisepticum* strain F over time after the organism was mixed with different amounts of MGA. A sample of MGA was initially diluted 1:10 in phosphate buffered saline (PBS). Then,  
30 the antiserum was further diluted by making 10 serial 1:2 dilutions by adding 0.5 ml of the previous dilution to 0.5 ml of PBS (dilutions 1:20 through 1:10240). One vial of *M. gallisepticum* F stock culture was thawed at room temperature and diluted 1:100. This 10(-2) stock solution contained  $5.8 \times 10(6)$  CFUs/ml. Bacterium-antibody complexes

- 33 -

were prepared by adding 0.4 ml of the  $5.8 \times 10^6$  stock solution to 0.4 ml of each of the 11 MGA dilutions. These complexes were allowed to react at room temperature for 30 minutes. Treatment 12 consisted of 0.4ml PBS added to 0.4 ml of the same bacterial stock solution.

- 5           Following the 30 minute reaction time, each of the 12 treatments was serially diluted in FMS from  $10^{-1}$  through  $10^{-8}$ . These tubes were incubated for 14 days and growth was determined at 41 hours, 47.5 hours, and at 14 days. (Growth of *M. gallisepticum* is detected in FMS by a color change. As bacterial growth increases the pH of the medium decreases, causing a color change in the pH indicator phenol red. As
- 10 growth occurs the color gradually changes from a deep red to orange and eventually to yellow. The degree of growth can be scored based on the medium color). Results are provided in Table 15.

Table 15

Growth of MgF after 0.4 ml of 12 antiserum dilutions, containing between .039 $\mu$ l and 40.0  $\mu$ l antiserum, were mixed with 0.4ml of an MgF stock solution

5

Treatment	MgF*	.4 ml MgF**	Growth after 41 Hours					Growth after 47.5 Hours					MgF Presence***
			10 (-1)	10 (-2)	10 (-3)	10 (-4)	10 (-5)	10 (-1)	10 (-2)	10 (-3)	10 (-4)	10 (-5)	
1	40 $\mu$ l	.4ml	-	-	-	-	-	-	+/-	-	-	-	10(-7)
2	20 $\mu$ l	.4ml	-	+/-	-	-	-	-	++	+	-	-	10(-6)
3	10 $\mu$ l	.4ml	+/-	+/-	+/-	-	-	-	++ +	+	-	-	10(-6)
4	5 $\mu$ l	.4ml	++	+/-	-	-	-	++ ++	++ +	+	-	-	10(-7)
5	2.5 $\mu$ l	.4ml	++ ++	+/-	-	-	-	++ ++	++ +	+/-	-	-	10(-6)
6	1.25 $\mu$ l	.4ml	++ ++	+/-	-	-	-	++ ++	++ +	+/-	-	-	10(-6)
7	.625 $\mu$ l	.4ml	++ ++	+/-	+/-	+/-	-	++ ++	++ +	-	-	-	10(-6)
8	.313 $\mu$ l	.4ml	++ ++	+	+/-	+/-	+/-	++ ++	++ +	+	+/-	+/-	10(-6)
9	.156 $\mu$ l	.4ml	++ ++	+/-	+/-	-	-	++ ++	++ +	+	-	-	10(-7)
10	.078 $\mu$ l	.4ml	++ ++	+/-	+/-	-	-	++ ++	++	+/-	-	-	10(-7)
11	.039 $\mu$ l	.4ml	++ ++	+/-	+/-	-	-	++ ++	++	+/-	-	-	10(-6)
12		.4ml	++ ++	+/-	-	-	-	++ ++	++	-	-	-	10(-7)

MgF\* = MgF antiserum in .4 ml

.4 ml MgF\*\* = .4 ml MgF 2.32x10(-6)

MgF Presence\*\*\* = Presence of MgF through dilution tube (after 14 days of incubation)

10

- (no growth; deep red)

+/- (growth just beginning; lighter red color than control tubes)

+

15

++ (moderate growth; deep orange)

+++ (moderate to heavy growth; light orange)

++++ (heavy growth; yellow)

Growth was detected in all 12 treatments by day 14 (Table 1). Growth was delayed in Groups 1-4, the groups containing the highest levels of MGA. Growth in these groups was not evident as early as in the groups with lower levels of MGA.

20

These findings suggest that the higher levels of MGA had a growth inhibiting effect on the bacterium.

#### EXAMPLE 16

##### 5 Growth Inhibition Effects of MGA on *M. gallisepticum*

A vial of *M. gallisepticum* strain F stock was thawed and diluted 1:100 in FMS. This 10(-2) dilution contained approximately  $5 \times 10^6$  CFUs/ml. One ml of the 10(-2) stock dilution was placed into each of eight dilution tubes after being thoroughly  
10 mixed. A certain amount of MGA was added to each tube (see Table 16) and the bacterium/antiserum complexes were mixed. They were incubated at room temperature for 15 minutes and then at 37degrees C. Growth was determined over the course of 13 days using color change in the FMS growth medium as an indicator. Results are given in Table 16.

15

Table 16

MgF 5.0 x 10(6) CFU/ml	µls of Mg Antiserum (MGA)	Growth after 27 hours at 37°C	Growth after 46 hours at 37°C	Growth after 13 days at 37°C
1.0 ml	No serum	+++	++++	++++
1.0 ml	48	no growth	no growth	++++
1.0 ml	24	no growth	no growth	++++
1.0 ml	12	no growth	no growth	++++
1.0 ml	6	no growth	+	++++
1.0 ml	3	no growth	++	++++
1.0 ml	1.5	no growth	+++	++++
1.0 ml	0.5	++	++++	++++

+ (light growth; light red)

++ (moderate growth; deep orange)

+++ (moderate to heavy growth; light orange)

20 +++++ (heavy growth; yellow)

Moderate growth of *M. gallisepticum* occurred within the first 27 hours of incubation in the tube that did not contain antiserum and growth was heavy in this tube by 46 hours at 37 degrees C (Table 16). Bacterial growth was not detected within the  
25 first 27 hours of incubation in the tubes that contained greater than or equal to 1.5µls MGA. After 46 hours of incubation, growth was still not detected in the tubes that contained 12, 24, and 48 µls of MGA. All tubes showed heavy growth of *M. gallisepticum* after 13 days at 37 degrees C. These results show that the bacterium was

- 36 -

present in all tubes, but that higher amounts of antiserum delayed growth for longer periods of time than did lesser amounts. The time that it took for growth to be detected appears to be directly proportional to the amount of MGA in the bacterium-antiserum complex.

5

**EXAMPLE 17****Effect of *M. gallisepticum*-MGA Complexes on Hatch**

10               Nine groups of eggs were inoculated at day 18 of incubation. Seven of the groups were inoculated with one of seven different *M. gallisepticum* strain F-MGA complexes. One group was inoculated with the bacterium only and another group was inoculated with a 1:4 dilution of FMS in PBS.

15               A vial of *M. gallisepticum* strain F stock was thawed and diluted 1:5 and 1:10 in 10% FMS and 90% PBS diluent. Appropriate amounts of these dilutions were used to create the bacterium-MGA complexes. Each egg received a 0.1 ml injection containing the same number of *M. gallisepticum* CFUs with the appropriate amount of antiserum for a particular group, except for Group 1. The eggs in Group 1 received 0.1 ml of the FMS/PBS diluent. The complexes were allowed to react together for 10 minutes before  
20               inoculation. The eggs were then incubated until hatch.

              The 1:10 dilution of the bacterial stock was titered by making 3 serial dilution series through 10(-9) and plating the 10(-6) dilutions of two dilution series TSA and incubating at 37°C. All tubes in all three serial dilution series showed *M. gallisepticum* growth. The titer was  $8 \times 10^8$  CFUs/ml. Hatchability results are provided in Table  
25               17.

Table 17

Group	MgF CFUs	Amount of Mg Antiserum (MGA)	#Eggs Hatched/ # Eggs Injected	# Healthy Chicks/ #Eggs Injected
1	0	0	13/14 (93%)	12/14 (86%)
2	$8.0 \times 10(6)$	40 $\mu$ l	10/11 (91%)	8/11 (73%)
3	$8.0 \times 10(6)$	20 $\mu$ l	10/10 (100%)	8/10 (80%)
4	$8.0 \times 10(6)$	10 $\mu$ l	6/11 (55%)	0/11 (0%)
5	$8.0 \times 10(6)$	5 $\mu$ l	10/10 (100%)	2/10 (20%)
6	$8.0 \times 10(6)$	2 $\mu$ l	6/11 (55%)	1/11 (9%)
7	$8.0 \times 10(6)$	0.5 $\mu$ l	7/11 (64%)	0/11 (0%)
8	$8.0 \times 10(6)$	.05 $\mu$ l	8/11 (73%)	0/11 (0%)
9	$8.0 \times 10(6)$	0	8/14 (57%)	0/14 (0%)

5

The MGA-*M.gallisepticum* complexes influenced the percent hatch and chick health. The groups that experienced hatches above 90% were the groups that contained the largest proportions of MGA to CFUs (with the exception of Group 4). The percentage of health chicks was much higher in Groups 2 and 3 than in other groups receiving MGA-bacterium complexes with less antiserum in the formulation. These findings indicate that certain MGA:bacterium ratios have the capability of protecting a developing chicken embryo by delaying and/or decreasing the pathogenic effects of the bacterium.

15

**EXAMPLE 18*****M. gallisepticum* strain F/ Antibody Complex Vaccine**

Six groups of 16 viable eggs were inoculated at day 18 of incubation. The 16 eggs in the negative control group (Group 6) were inoculated on day 18 with 0.1ml of diluent (1 part FMS in 9 parts PBS), and hatched in a large hatcher unit that contained no other eggs.

A vial of *M. gallisepticum* strain F stock was thawed at room temperature and diluted 1:5 (stock 2). The titration of stock 2 showed that it contained  $7 \times 10(7)$  CFUs/ml. Stock 2 was then divided into 0.9 ml aliquots and combined with 0.5, 10, 20 or 40  $\mu$ l of MGA. Once mixed, the bacterium-MGA formulations were allowed to incubate at room temperature for 15 minutes. These bacterium-MGA complexes were administered to eggs of each group in 0.1 ml doses. Group 1 eggs received inoculations

25

- 38 -

containing only bacteria. Each group of 16 eggs was then placed in separate small hatcher units until day of hatch. The MGA-*M. gallisepticum* formulations tested CFUs are shown in Table 18.

5 The remaining stock 2 dilution was titrated in three separate serial 10-fold dilutions to 10(-9) using FMS. The 10(-4), 10(-5) and 10(-6) dilution tubes in each series was plated in quadruplicate on FMS agar and incubated at 37°C for 9 days.

On the day of hatch, Groups 1-5 were processed. Normal, healthy looking chicks were sampled for the presence of *M. gallisepticum* by swabbing the choanal cleft with a sterile swab and inoculating tubes containing 1.8 mls of FMS. After the chicks  
10 were processed, the sampled chicks from each group were placed in a P2 containment room. Each group was placed in a separate brooder cage and no two cages were in contact.

The chicks in vehicle control Group 6 experienced a delayed hatch and were processed the day following the hatch of groups 1-5. Ten control birds were swabbed  
15 for the presence of *M. gallisepticum* and were placed in a brooder cage in a separate P2 containment room. On 21 days of age, all surviving chicks were bled and serum collected for determination of antibodies to *M. gallisepticum* by serum plate agglutination (SPA) and ELISA. Results are provided in Table 18.



Table 18

Group	1*	2	3	4	5	6
#CFUs/egg	1.4x10(6)	3.5x10(6)	3.5x10(6)	3.5x10(6)	3.5x10(6)	no CFUs
MGA/egg	0µl	5µl	10µl	20µl	40µl	100µl diluent
# Normal Hatched	0	0	0	8	12	10
# Clinically Affected Hatched	9	7	8	3	2	1
Unhatched/Dead	7	9	8	5	2	5
% Hatched	56.25	43.75	50.0	68.75	87.5	68.75
% Normal Hatched	0	0	0	50	75	62.5
# Chicks Placed	0	0	3	9	10	10
# Chicks Alive at 3 weeks	NA	NA	1	2	7	10
MgF Reisolation # Positive/ # Sampled	Not Tested	3/3	3/3	9/9	9/10	0/10
SPA** # Positive/ # Sampled	NA	NA	1/1	2/2	6/7	0/10
Mean ELISA Titer	NA	NA	599	643	645	0

\* Due to a calculation error, the eggs in Group 1 received 1.4x10(6) CFUs.

5 \*\* All positive SPA reactions were scored a 3 or higher on a scale of 0-4, with 4 being the strongest reaction.

The eggs receiving the bacterium without antiserum (Group 1) and the two lower amounts of MGA plus *M. gallisepticum* (Groups 2 and 3) had the lowest percent hatches and no normal birds hatched. The vehicle control group (Group 6) experienced a delayed hatch as well as a poorer hatch than expected. This probably was caused by the fact that the eggs were incubated in a large hatcher intended for the incubation of 2000 eggs. Despite this hatch problem, 10 chicks were healthy and they tested negative for *M. gallisepticum* isolation and the two serum antibody tests. Groups 4 and 5 experienced percent hatches and percent normal hatches that showed great improvement over those of Groups 1, 2 and 3. The eggs in these two groups received higher levels of MGA and Group 5 (3.5x10(6)CFU +40µl MGA) experienced the best hatch of all the groups. All birds remaining alive at the time of serum collection were

- 40 -

healthy. The antibody response to *M. gallisepticum* measured by the SPA test and ELISA indicate that the *M. gallisepticum* strain F complex vaccines were efficacious for birds in Groups 3, 4 and 5.

- It is interesting to note that one bird in Group 5 was negative for *M. gallisepticum* reisolation at hatch, the SPA test, and ELISA. All the other birds that were sampled from Groups 3, 4 and 5 tested positive in each case. It appears that one bird in Group 5 never became infected.

- Examples 14-18 were designed to test the usefulness of a bacteria:antibody vaccine complex. The data support the concept that addition of specific antiserum (specific for the vaccine bacteria) to live bacteria in the appropriate ratio provides protection to the chick embryo by decreasing or delaying the pathogenic effects of the bacterium while at the same time allowing an efficacious immune response to develop in the hatchlings, as evidenced by an active humoral immune response.

- 41 -

**THAT WHICH IS CLAIMED IS:**

1. A method of producing active immunity against a bacterial disease in a subject, said method comprising:
  - administering to the subject a vaccine conjugate comprising a live bacteria and a neutralizing factor bound to the live bacteria;
  - 5 the neutralizing factor selected from the group consisting of antibodies and antibody fragments;
  - the antibody or antibody fragment capable of neutralizing the live bacteria;
  - the vaccine conjugate administered in an amount effective to produce an
  - 10 immune response to the live bacteria in the subject.
2. A method according to Claim 1, wherein the live bacteria is capable of causing disease in the subject.
- 15 3. A method according to Claim 1, wherein the neutralizing factor is selected from the group consisting of IgG immunoglobulins and IgG immunoglobulin fragments.
4. A method according to Claim 1, wherein the subject is a bird and
- 20 said administering step is carried out in ovo.
5. A vaccine preparation useful for producing active immunity against a bacterial disease in a subject, said vaccine preparation comprising:
  - a pharmaceutically acceptable formulation comprising a vaccine
  - 25 conjugate;
  - said vaccine conjugate comprising a live bacteria and a neutralizing factor bound to said live bacteria;
  - said neutralizing factor selected from the group consisting of antibodies and antibody fragments;
  - 30 said antibody or antibody fragment capable of neutralizing said

- 42 -

live bacteria; and  
said vaccine conjugate included in said pharmaceutically acceptable  
formulation in an amount effective to produce an immune response to said live bacteria in  
said subject.

5

6. A vaccine preparation as claimed in Claim 5, wherein said live  
bacteria is capable of causing disease in said subject.

7. A vaccine preparation as claimed in Claim 5, wherein said  
10 pharmaceutically acceptable formulation is lyophilized.

8. A method of producing active immunity against a protozoal  
disease in a subject, said method comprising:  
administering to the subject a vaccine conjugate comprising a live  
15 protozoa and a neutralizing factor bound to the live protozoa;  
the neutralizing factor selected from the group consisting of  
antibodies and antibody fragments;  
the antibody or antibody fragment capable of neutralizing the live  
protozoa;  
20 the vaccine conjugate administered in an amount effective to produce an  
immune response to the live protozoa in the subject.

9. A method according to Claim 8, wherein the live protozoa is  
capable of causing disease in the subject.

25

10. A method according to Claim 8, wherein the live protozoa is an  
avian protozoa and the subject is a bird.

11. A method according to Claim 10, wherein the live protozoa is an  
30 *Eimeria* species.

12. A method according to Claim 10, wherein the live protozoa is *E.*

*tenella* or *E. acervulina*.

13. A method according to Claim 8, wherein the neutralizing factor is selected from the group consisting of IgG immunoglobulins and IgG immunoglobulin fragments.

5

14. A method according to Claim 8, wherein the neutralizing factor is of polyclonal origin.

15. A method according to Claim 8, wherein the neutralizing factor is  
10 of monoclonal origin.

16. A method according to Claim 8, wherein the subject is administered the vaccine conjugate by a method selected from the group consisting of subcutaneous administration, intraperitoneal administration, and intramuscular  
15 administration.

17. A method according to Claim 8, wherein the subject is a bird and said administering step is carried out in ovo.

20 18. A vaccine preparation useful for producing active immunity against a protozoal disease in a subject, said vaccine preparation comprising:

a pharmaceutically acceptable formulation comprising a vaccine conjugate;

25 said vaccine conjugate comprising a live protozoa and a neutralizing factor bound to said live protozoa;

said neutralizing factor selected from the group consisting of antibodies and antibody fragments;

said antibody or antibody fragment capable of neutralizing said live protozoa; and

30 said vaccine conjugate included in said pharmaceutically acceptable formulation in an amount effective to produce an immune response to said live protozoa in said subject.

- 44 -

19. A vaccine preparation as claimed in Claim 22, wherein said live protozoa is capable of causing disease in said subject.

5           20. A vaccine preparation as claimed in Claim 22, wherein said pharmaceutically acceptable formulation is lyophilized.

          21. A vaccine preparation as claimed in Claim 22, wherein said live protozoa is an avian protozoa.

10

          22. A vaccine preparation as claimed in Claim 22, wherein said live protozoa is an *Eimeria* species.

          23. A vaccine preparation as claimed in Claim 22, wherein said live protozoa is *E. tenella* or *E. acervulina*.

15

          24. A method of producing active immunity against a mycoplasma in a subject, comprising:

                  administering to the subject a vaccine conjugate comprising a live mycoplasma and a neutralizing factor bound to said mycoplasma, where the neutralizing factor is selected from the group consisting of antibodies and antibody fragments, said antibody or antibody fragment capable of neutralizing said live mycoplasma;

                  the vaccine conjugate administered in an amount effective to produce an immune response to the live mycoplasma in the subject.

25

          25. A method according to claim 24 wherein said mycoplasma is *Mycoplasma gallisepticum*.

30           26. A method according to Claim 24, wherein the neutralizing factor is selected from the group consisting of IgG immunoglobulins and IgG immunoglobulin fragments.

-45-

27. A method according to Claim 24, wherein the subject is a bird and said administering step is carried out *in ovo*.

1/3

FIG. 1






FIG. 2





2/3

FIG. 3



3/3

FIG. 4





FIG. 5



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/17259

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K39/40 A61K39/395 //(A61K39/40,39:02),(A61K39/395,39:002)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 91 04749 A (EMBREX INC) 18 April 1991 see the whole document ---	1-27
Y	WO 93 01832 A (LIOTET SERGE ;CATALAN FRANCOIS (FR); GOUCHET FRANCK (FR)) 4 February 1993 see the whole document ---	1-27
A	GB 1 105 136 A (HULLINGER LOTTE) 25 March 1964 see the whole document ---	1-27
P,X	WO 96 40233 A (PFIZER ;EVANS NIGEL A (US); FINDLY R CRAIG (US); WEBER FREDERICK H) 19 December 1996 see the whole document -----	8-23

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

29 January 1998

Date of mailing of the international search report

13. 02. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Müller, F

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 97/17259

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-4, 8-17, 24-27  
because they relate to subject matter not required to be searched by this Authority, namely:  
although claims 1-4, 8-17, 24-27 are directed to a method of treatment (or diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. .nal Application No

PCT/US 97/17259

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9104749 A	18-04-91	AT 150976 T AU 6714290 A AU 641958 B CA 2066661 A DE 69030383 D DE 69030383 T EP 0495008 A EP 0755684 A ES 2102368 T IL 95867 A JP 5501565 T US 5397569 A US 5397568 A W0 9104750 A	15-04-97 28-04-91 07-10-93 03-04-91 07-05-97 23-10-97 22-07-92 29-01-97 01-08-97 07-10-94 25-03-93 14-03-95 14-03-95 18-04-91
W0 9301832 A	04-02-93	CH 682455 A AT 134879 T CA 2092302 A DE 69208854 D DE 69208854 T EP 0550724 A ES 2085025 T	30-09-93 15-03-96 26-01-93 11-04-96 10-10-96 14-07-93 16-05-96
GB 1105136 A		NONE	
W0 9640233 A	19-12-96	AU 5478396 A BG 100641 A CN 1145167 A CZ 9601642 A HU 9601553 A PL 314640 A SI 9600182 A SK 72796 A	19-12-96 31-07-97 19-03-97 12-03-97 28-03-97 09-12-96 30-06-97 06-08-97

